

POSTER PRESENTATION

Open Access

Surveillance of central-line associated bloodstream infections in ICU at a Malaysian medical centre

A Sulong^{1*}, N Awang Jalil¹, R Ramli¹, M Mohd Yusoff²

From International Conference on Prevention & Infection Control (ICPIC 2011)
Geneva, Switzerland. 29 June – 2 July 2011

Introduction / objectives

We conducted a prospective surveillance study to determine the incidence and epidemiology of CLABSI in the intensive care unit (ICU) at a tertiary teaching medical centre using CDC definition of a bloodstream infection that develops in a patient who had a central line at the time of, or within 48 hours before, the onset of the infection.

Methods

A cross section observational study was conducted in a 24-bed GICU over 10-months period from April 2008 until January 2009 among patients with central venous catheters (CVCs) inserted in the ICU or operation theatre.

Results

A total of 20 CLABSI cases were identified among 155 CVCs in 100 patients with 3106 catheter days. The overall rate of CLABSI was 6.4 per 1000 catheter-days and device-utilization (DU) ratio of 0.81. The mean length of ICU stay for CLABSI and non-CLABSI cases was 37.2 days and 17.4 days respectively, while the median length of ICU stay for CLABSI cases was 16.0 days and for non-CLABSI cases was 10.0 days. This contributed to 6 extra days of ICU stay in CLABSI cases. CVCs inserted via the femoral vein were associated with higher infection rate of 22.2% followed by those of internal jugular vein (15.4%) and subclavian vein (5.1%). Gram-negative bacteria accounted for 50% of the CLABSI cases whereas gram-positive cocci and fungi caused 35% and 15% of these infections respectively.

Conclusion

Both of the CLABSI rate of 6.4 per 1000 catheter-days and DU ratio of 0.81 were above the 90th percentile of the NHSN benchmark. Comparing our findings to the INNIC 2003-2008 studies with 7.4 CLABSI per 1000 catheter-days, our CLABSI rate was lower. However, our DU ratio was higher. Hence, interventions aimed at improving outcomes related to CVCs should seriously be considered.

Disclosure of interest

None declared.

Author details

¹Dept of Medical Microbiology & Immunology, Faculty of Medicine, Universiti Kebangsaan Malaysia Medical Centre, Kuala Lumpur, Malaysia.

²Dept of Anaesthesiology, Faculty of Medicine, Universiti Kebangsaan Malaysia Medical Centre, Kuala Lumpur, Malaysia.

Published: 29 June 2011

doi:10.1186/1753-6561-5-S6-P212

Cite this article as: Sulong et al.: Surveillance of central-line associated bloodstream infections in ICU at a Malaysian medical centre. *BMC Proceedings* 2011 5(Suppl 6):P212.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Dept of Medical Microbiology & Immunology, Faculty of Medicine, Universiti Kebangsaan Malaysia Medical Centre, Kuala Lumpur, Malaysia
Full list of author information is available at the end of the article